

Product:	Anti-17β-Nortestosterone Antibody					
Synonyms:	Not available					
Immunogen:	17β-19-Nortestosterone-BTG					
	1 mg	PAS9708	Host Species:	Ovis aries (Sheep)	Isotype:	IgG
Pack sizes: Cat. No.	10 mg	PAS9557	Host Breed:	Texel	Format:	Clear liquid
			Clonality:	Polyclonal	рН:	7.4
IgG Concentration:			Buffer:			
Lot dependent. Determined @ 280nm.		20mM Phosphate, 150mM Sodium Chloride.				
Recommended Working Concentration*:			Preservative:			
10 μg/mL			0.09% Sodium Azide.			
*The application notes inc dilutions/concentrations sh	lude recommen nould be determ	ded starting dilutions; optimal ined by the end user.				
Assessment Method:			Species Reactivity:			
Competitive ELISA.			N/A			
Method of Purification:			Storage:			
Salt fractionation.			Can be stored for up to 3 months at +2°C - +8°C. For long term storage, aliquot and store at ≤-20 °C. Avoid repeated freeze/thaw cycles. Product should be protected from light exposure.			
Recommende	d Applica	ations:	<u> </u>			
		for the development of im	munoassays or imr	nunoaffinity purification	columns.	
Sensitivity:			Target Specificity*:			
10 ng/mL 17Beta-19Nortestosterone produces 81% inhibition in a competitive ELISA employing 17β-Nortestosterone polyclonal antibody.			Data not availabl	e.		
Related Produ	cts:					
17β-Nortestost	erone-HR	RP, HRP9369, Kit size: 0.5	mL.			

Notes & Precautions:

Antibody can be affinity purified on request. This product as supplied is intended for research applications only, not for use in therapeutic or diagnostic applications without the expressed written authorization of Randox BioReagents. A safety data sheet (SDS) can be supplied upon request. Vial should be centrifuged briefly before opening to ensure all material is removed from the vial cap.

28-Apr-22 Information correct at time of going to print

