

Discovery | Drug Development | Diagnostics

Product:	Anti-JWH-018 (SPICE) Antibody						
Synonyms:	Not available						
Immunogen:	JWH-018-BTG						
	1 mg	PAS10172	Host Species:	Ovis aries (Sheep)	Isotype:	lgG	
Pack sizes: Cat. No.	10 mg	PAS10173	Host Breed:	Texel	Format:	Clear liquid	
			Clonality:	Polyclonal	рН:	7.4	
IgG Concentration: Lot dependent. Determined @ 280nm.			Buffer: 20mM Phosphate	Buffer: 20mM Phosphate, 150mM Sodium Chloride.			
Recommended Working Concentration*: 5 µg/mL *The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.			Preservative: 0.09% Sodium Az	Preservative: 0.09% Sodium Azide.			
Assessment Method:			Species Reactivi	Species Reactivity:			
Competitive ELISA.			N/A	N/A			
Method of Purification:			Storage:	Storage:			
Salt fractionation.			For long term stor Avoid repeated fre	Can be stored for up to 3 months at +2°C - +8°C. For long term storage, aliquot and store at ≤-20 °C. Avoid repeated freeze/thaw cycles. Product should be protected from light exposure.			
Recommended	Applica	itions:					
The antibody is suitable for the development of immunoassays or immunoaffinity purification columns.							

Sensitivity:

10 ng/mL JWH018 produces 71% inhibition in a competitive ELISA employing JWH-018 (SPICE) polyclonal antibody.

Target Specificity*:

ELISA microtitre plate based and expressed as % crossreactivity*. The antibody exhibited crossreactivity for the following compounds: Psilocin <0.03%, Indole-3-carboxylic acid <0.03%, 5-Hydroxyindole-3-acetic acid <0.03% and 1-Naphthoic acid <0.03%.

* Cross reactivity profile may vary with tracer used

Related Products:

JWH-018 (SPICE)-HRP, HRP9523, Kit size: 0.5 mL.

Notes & Precautions:

Antibody can be affinity purified on request. This product as supplied is intended for research applications only, not for use in therapeutic or diagnostic applications without the expressed written authorization of Randox BioReagents. A safety data sheet (SDS) can be supplied upon request. Vial should be centrifuged briefly before opening to ensure all material is removed from the vial cap.

28-Apr-22

Information correct at time of going to print

