

Product:	Anti-Theophylline Antibody						
Synonyms:	Not available						
Immunogen:	Theophylline-BTG						
	1 mg	PAS10196	Host Species:	Ovis aries (Sheep)	Isotype:	IgG	
Pack sizes: Cat. No.	10 mg	PAS10197	Host Breed:	Texel	Format:	Clear liquid	
			Clonality:	Polyclonal	pH:	7.4	
IgG Concentration:			Buffer:				
Lot dependent. Determined @ 280nm.			20mM Phosphate, 150mM Sodium Chloride.				
Recommended Working Concentration*:  5 µg/mL  *The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.			Preservative: 0.09% Sodium Azide.				
Assessment Method:			Species Reactivity:				
Competitive ELISA.			N/A				
Method of Purification:			Storage:				
Salt fractionation.			Can be stored for up to 3 months at +2°C - +8°C. For long term storage, aliquot and store at ≤-20 °C. Avoid repeated freeze/thaw cycles. Product should be protected from light exposure.				
Recommende	d Applica	ations:	1				
The antibody is	suitable	for the development of im	munoassays or imr	munoaffinity purification	columns.		
Sensitivity:			Target Specificity*:				
100 ng/mL Theophylline produces 93% inhibition in a competitive ELISA employing Theophylline polyclonal antibody.			ELISA microtitre plate based and expressed as % crossreactivity*. The antibody exhibited crossreactivity for the following compounds: Aminophylline 59.5%, 8-Chlorotheophylline 13.4%, Caffeine <1.7%, Theobromine <1.7%, 1,3-Dimethyluric acid 11.0%, 1,7-Dimethylxanthine <1.7%, 1-Methyluric acid <1.7% and 3-Methylxanthine <1.7%.				
			* Cross reactivity profile may	* Cross reactivity profile may vary with tracer used.			
Related Produ	icts:						

Theophylline-HRP, HRP9537, Kit size: 0.5 mL.

## Notes & Precautions:

Antibody can be affinity purified on request. This product as supplied is intended for research applications only, not for use in therapeutic or diagnostic applications without the expressed written authorization of Randox BioReagents. A safety data sheet (SDS) can be supplied upon request. Vial should be centrifuged briefly before opening to ensure all material is removed from the vial cap.

12-May-22

Information correct at time of going to print

