

Interleukin-3 Human, Recombinant, *E. coli*

Cat. No.: RCP9000 **Size**: 10μg

Synonym: IL-3; Multi-potential Colony Stimulating Factor (MCSF);

Hematopoietic Growth Factor; P-Cell Stimulating Factor; Growth Factor (MCGF); Multi-lineage Colony Stimulating

Mast Cell Factor.

Description:

Interleukin-3 is a pleiotropic cytokine produced primarily by activated T cells. IL-13 is thought to function via specific cell surface receptors to stimulate the proliferation, differentiation and survival of haematopoietic cell lines. IL-3 has also been shown to affect the functional activity of a variety of other cell types including mast cells, eosinophils, megakaryocytes and basophils

megakaryocytes and basophils

RANDOX recombinant Interleukin-3 comprises a 133 amino acid fragment (20-152) corresponding to the mature Interleukin-3 protein and is expressed in *E. coli* with an amino-terminal hexahistidine tag. This product is for research use only and is not intended for diagnostic or the reposition use.

therapeutic use.

Form: Liquid.

Supplied in 1x Laemmli Buffer (25mM Tris-HCl pH6.8, 50mM

ווט,

1% (w/v) SDS, 0.1% (w/v) Bromophenol Blue, 2.5% Glycerol).

Purity: Single band on Western blot.

References: Lindemann A. & Mertelsmann R. (1993) Cancer Invest.

11(5): 609-623.

