

Interleukin-8 Human, Recombinant, E. coli

Cat. No.: RCP9276 Size: **10μg**

RCP9277 100μg RCP9125 1000µg

IL-8, CXCL8, Monocyte-derived neutrophil chemotactic Synonym:

> factor, MDNCF, T-cell chemotactic factor, Neutrophilactivating protein 1, NAP-1, Protein 3-10C, Granulocyte chemotactic protein 1, GCP-1, Monocyte derived

neutrophil activating peptide, MONAP, Emoctakin.

Description: Interleukin-8 (IL-8) belongs to the neutrophil-specific CXC

family of chemokines. It is one of the initial cytokines released from a variety of cell types, including T cells, endothelial cells and fibroblasts, in response to an inflammatory stimulus and acts by recruiting neutrophils, T-cells and basophils to the site of inflammation. Elevated Interleukin-8 levels are associated with the onset of a

variety of disease states.

RANDOX recombinant IL-8 comprises the 77 amino acid fragment (23-99) corresponding to the mature IL-8 protein and is expressed in E. coli with an amino-terminal hexahistidine tag. This product is for research use only and is not intended for diagnostic or therapeutic use.




50.7 kDa

114 kDa 88 kDa

Purity: >95% by SDS-PAGE 35.5 kDa

22.0 kDa



References:

Baggiolini, M & Clark-Lewis I (1992) FEBS Letters 307 (1):

97-101.

