

VCAM-1 Fragment Human, Recombinant, *E. coli*

Cat. No.: RCP9075 **Size**: 10μg

Synonym: Vascular cell adhesion molecule 1, INCAM-100, Vascular

cell adhesion protein 1 precursor, CD-106 antigen.

Description: VCAM-1 is a cell surface sialoglycoprotein expressed by

cytokine activated endothelium. The protein has a number of functions including the regulation of leukocyte migration, leukocyte-endothelial cell adhesion and signal transduction. VCAM-1 may also play a role in a number of

inflammatory diseases.

RANDOX recombinant VCAM fragment comprises a 203 amino acid fragment (109-331) corresponding to the Ig-like C2 type domains 2 and 3 and is expressed in *E. coli* with an amino-terminal hexahistidine tag. This product is for research use only and is not intended for diagnostic or

therapeutic use.

Form: Liquid.

Supplied in 1x Laemmli Buffer (25mM Tris-HCl pH6.8, 50mM DTT, 1% (w/v) SDS, 0.1% (w/v) Bromophenol Blue,

2.5% Glycerol).

Purity: Single band on Western blot.

References: Steeber, D. A & Tedder, T.F. (2000) Immuno. Res. 22(2-

3): 299- 317.

Albelda, S.M et al. (1994) FASEB J. 8(8): 504-512.

