

VEGFR-1

Human, Recombinant, *E. coli*

| | | | |
|------------------|---------|--------------|--------|
| Cat. No.: | RCP9016 | Size: | 10µg |
| | RCP9017 | | 100µg |
| | RCP9138 | | 1000µg |


Synonym: Vascular endothelial growth factor receptor-1 [precursor]; VEGFR-1; EC 2.7.1.112; Vascular permeability factor receptor ; Tyrosine-protein kinase receptor FLT; FLT-1; Tyrosine-protein kinase FRT; FMS-like tyrosine kinase 1.

Description: VEGFR-1 belongs to a family of five receptor tyrosine kinase receptors that are specific to a number of angiogenic factors including VEGF and PDGF, and are therefore thought to play a role in the angiogenic process. VEGFR-1 is expressed in two forms, predominantly on endothelial cells, giving rise to a full-length membrane bound receptor and a truncated soluble receptor. Elevated levels of the soluble form of this VEGF receptor may be used in the diagnosis of pathological conditions.

RANDOX recombinant VEGFR-1 comprises a 298 amino acid fragment (31-328) corresponding to the IgG-like domains 1-3 from the mature soluble VEGFR protein. VEGFR-1 is expressed in *E. coli* with an amino-terminal hexahistidine tag. This product is for research use only and is not intended for diagnostic or therapeutic use.

Form: Liquid

Purity: >95% by SDS-PAGE



Reference(s): Starovasnik, M. A *et al.* (1999) *J. Mol. Biol.* **296 (3)** : 531-544.

Kanno, S. *et al.* (2000) *Oncogene* **19 (17)** : 2138-2146.