

sCD40L (113-261) Human, Recombinant, *E. coli*

Cat. No.: RCP9427 **Size:** 10μg

RCP9428 100μg RCP9429 1000μg

Synonym: CD40-L, TNF-related activation protein, Tumour Necrosis

Factor Ligand Superfamily member 5, TRAP, T cell antigen Gp39, CD154 antigen, sCD40, IGM, IMD3,

HIGM1, T-BAM, TNFSF5, hCD40L.

Description: sCD40L is contained in <u>platelet</u> granules and thus its

presence in the blood is a marker of <u>platelet</u> activation. By interacting with <u>CD40</u>, which is found on endothelial and smooth muscle cells, sCD40L may trigger the release of inflammatory mediators, lead to increased activity of <u>matrix</u> metalloproteinases and activation of the coagulation

cascade.


RANDOX recombinant sCD40L comprises a 149 amino acid fragment (113-261) corresponding to the receptor binding domain of the sCD40L protein and is expressed in *E. coli* with an amino-terminal hexahistidine tag. This product is for research use only and is not intended for

148kDa



Form: Liquid

Purity: >95% by SDS-PAGE



References: Schönbeck, U. et al. (2001) <u>Cellular</u> and Molecular Life Sciences **58 (1):** 4-43.

Lineberry, N.B. et al. (2008). Journal of Immunology **181** (3):1622-6.

